

Tertiary Eclogite Facies Metamorphism in the Greater Himalayan Sequence: Evidence from Zircon U-Pb Geochronology and Trace Element Geochemistry

Djordje Grujic¹, Clare Warren², Joseph L. Wooden³

¹ Department of Earth Sciences, Dalhousie University, Halifax, NS B3H 4J1, Canada, dgrujic@dal.ca

² Earth and Environmental Sciences, The Open University, Milton Keynes MK7 6AA, England

³ Department of Geological and Environmental Science, Stanford University, Stanford, CA 94305, U.S.A.

Mafic and pelitic granulites exposed in the Greater Himalayan Sequence (GHS) in the eastern Himalayan kingdom of Bhutan preserve textural evidence for a precursor high-pressure metamorphic event, the precise conditions of which are generally unrecoverable due to the later high-temperature overprint (Groppo and others, 2007). As high pressure metamorphism is rare in the Himalayas, especially in the eastern parts of the orogen, their thermobarometrical and geochronological evolution place important constraints on the geodynamic evolution of the Himalaya in particular, and continental collisions in general. We report SHRIMP-RG trace element (REE) and U–Pb zircon geochronological data, collected by the same instrument and on adjacent spots of the same crystal. These data suggest that zircons crystallized at 14–15 Ma over a temperature range of ca. 705–815 °C. This age is interpreted to indicate the timing of HP metamorphism due to the lack of negative Eu anomaly, the depleted heavy REE signature and the temperatures of crystallization.

U–Th–Pb Monazite ages indicate that the near-peak T conditions in this area were attained shortly after the somewhat lower grade area underneath separated by a ductile shear zone. Ti-in-zircon and Zr-in-rutile geothermometry further help to establish links between accessory mineral crystallization and metamorphism. Finally, crystallization ages of deformed leucogranites suggest a concomitant shift of deformation along the roof normal-geometry shear zone towards the interior of the orogen (Kellett and others, 2009). These data are consistent with multiple pulses of GHS exhumation in Bhutan from south to north (Hollister and Grujic, 2006), possibly by tectonic forcing over an incoming Indian crustal ramp.

We suggest that rocks in GHS in the eastern Himalaya were buried to greater depths and subjected to greater temperatures than in the central parts of the orogen, and were exhumed rapidly during the later stages of orogenic evolution. The formation of the Himalaya is currently explained by two contrasting tectonic models that differ in their predictions for the sequence of deformation along the main structures. We conclude that deformation and metamorphism distribution in the Bhutan Himalaya is compatible with the predictions of geodynamic models of Himalaya applying concepts of the channel flow hypothesis (Jamieson and others, 2006).

References

- Hollister, L. S. and Grujic, D., 2006, Pulsed channel flow in Bhutan, in: Law, R. D., Searle, M. P. & Godin, L. (eds) Channel Flow, Extrusion, and Exhumation in Continental Collision Zones, Geological Society, London, Special Publications, 268, 415–423.
- Groppo, C., Lombardo, B., Rolfo, F. and Pertusati, P., 2007, Clockwise exhumation path of granulitized eclogites from the Ama Drime range (Eastern Himalayas), *Journal of Metamorphic Geology*, 25, 51–75.
- Jamieson, R.A., Beaumont, C., Nguyen, M.H. and Grujic, D., 2006, Provenance of the Greater Himalayan Sequence and associated rocks: Predictions of channel flow models, in: Law, R. D., Searle, M. P. & Godin, L. (eds) Channel Flow, Extrusion, and Exhumation in Continental Collision Zones, Geological Society, London, Special Publications, 268, 165–182.
- Kellett, D., Grujic, D. and Erdman, S., 2009, Miocene structural reorganization of the South Tibetan detachment, eastern Himalaya: Implications for continental collision, *Lithosphere*, 1, 259–281.